

## **Ichnofacies of the Gething Formation of NE British Columbia**

Murray K. Gingras\*

Department of Earth and Atmospheric Sciences  
1-26 Earth Sciences Building, University of Alberta, Edmonton, AB T6G 2E3  
mgingras@ualberta.ca

and

John Paul Zonneveld

Geological Survey of Canada, Calgary, AB

### **ABSTRACT**

The Gething Formation in northeast British Columbia is a variably thick, heterolithic succession of continental and marginal-marine facies. This unit was deposited on the south Flank of the Keg River Highlands in the northern Western Canada Foreland Basin (~58°N, 120°W). Although Gething sandstone bodies are exploration targets further south, it is not considered an exploration target in the study area: due largely to its complex and laterally discontinuous nature. However, in the studied interval several of the lithofacies possess good reservoir properties.

New subsurface research in the area of Ring and Kahntah River gas fields shows that the Gething is represented by three valleys that debouch southward into a large structural basin. In their northernmost (upstream) reaches, the Gething Formation strata are dominated by unbioturbated, high-angle cross-bedded and current-ripple laminated upper-fine-grained sandstones, rich in organic detritus. These are interpreted as fluvial / bayhead delta deposits.

The continental lithofacies grade southwards into bioturbated, low- and high-angle cross-bedded, lower to upper-fine-grained sandstones with thin, bioturbated siltstone and mudstone intervals. Rhythmic bedding, mud flasers, and sedimentary couplets, and the trace fossils *Planolites*, *Skolithos*, *Cylindrichnus*, *Arenicolites*, *Teichichnus*, and fugichnia are common. These strata may represent brackish-water bay deposits. South of this, the Gething is dominated by interstratified, sporadically bioturbated sand, silt and mud, sedimentologically similar to the aforementioned brackish channel deposits, but with a more diverse and robust ichnofauna (the previous assemblage as well as *Thalassinoides*, rare *Helminthopsis* and *Phycosiphon*). These deposits are ascribed to tidal channels in the larger, more marine bay.