Petroleum Potential of the Middle-Upper Devonian Canol Formation in North Yukon: Report on Field Investigations in 2013

Tiffani A. Fraser¹ and Matt Hutchison¹

Abstract

The Yukon Geological Survey is conducting research to evaluate the petroleum potential of Devonian-Carboniferous shale units in north Yukon. The Middle-Upper Devonian Canol Formation was the focus of field studies in summer 2013. Stratigraphic sections in the southern Richardson and northern Ogilvie mountains were systematically measured and sampled in an effort to characterize and correlate the Canol Formation within the Yukon, and to assess the formation's regional hydrocarbon potential, as both a source rock and a shale reservoir. This study presents preliminary results for three Canol Formation stratigraphic sections. The information acquired from these outcrops will be instrumental in understanding the Canol Formation in the subsurface of Eagle Plain Basin and Peel Plateau and Plain oil and gas regions. The study is timely as the Canol Formation is currently being explored as an unconventional hydrocarbon resource in the Northwest Territories' central Mackenzie Valley.

References Cited

Fraser, T.A. and Hogue, B., 2007. List of wells and formation tops, Yukon Territory, version 1.0. Yukon Geological Survey Open File 2007-05.

Fraser, T.A., Allen, T.L., Lane, L.S. and Reyes, J.C., 2012. Shale gas potential of Devonian shale in north Yukon: Results from a diamond drillhole study in western Richardson Mountains. In Yukon Exploration and Geology 2011, edited by K.E. MacFarlane and P.J. Sack, Yukon Geological Survey, p. 45-74.

Gal, L.P., Pyle, L.J., Hadlari, T., and Allen, T.L., 2009. Chapter 6 – Lower to Upper Devonian Strata, Arnica-Landry Play, and Kee Scarp Play. In Regional Geoscience Studies and Petroleum Potential, Peel Plateau and Plain, Northwest Territories and Yukon: Project Volume, edited by L.J. Pyle and A.L. Jones, Northwest Territories Geoscience office and Yukon Geological Survey, NWT Open File 2009-02 and YGS Open File 2009-25, p. 187-289.

Hayes, B.J.R., and Archibald, H.B., 2012. Scoping study of unconventional oil and gas potential, Yukon. Yukon Geological Survey Miscellaneous Report 7.

Lenz, A.C. 1972. Ordovician to Devonian history of northern Yukon and adjacent District of Mackenzie. Bulletin of Canadian Petroleum Geology, vol. 20, p. 321-361.

¹Yukon Geological Survey, Whitehorse, Yukon, Canada

Link, C.M., Bustin, R.M. and Snowdon, L.R., 1989. Petroleum source potential and depositional setting of Phanerozoic strata in northern Yukon and northwestern district of Mackenzie. Bulletin of Canadian Petroleum Geology, vol., 37, no. 3, p. 293-315.

Mathews, W.H. and Bustin, R.M., 1984. Why do the smoking Hills smoke? Canadian Journal of Earth Sciences, 21, pp. 737-742.

Northern Cross Yukon Limited, 2013. Eagle Plain 3D seismic and exploration project review. Accessed on the internet at http://www.northerncrossyukon.ca/upload/news_item/9/01/3d-seismic-exploration-project-review.pdf, January 9, 2014.

Osadetz, K, G., MacLean, B.C., Morrow, D.W., Dixon, J. and Hannigan, P.K., 2005. Petroleum resource assessment, Peel Plateau and Plain, Yukon Territory, Canada. Yukon Geological Survey Open File2005-03, Geological Survey of Canada Open File 4841.

Pigage, L.C., 2009. Yukon Table of Formations v 3.2. Yukon Geological Survey and Oil and Gas Resources Branch.

Pugh, D.C. 1983. Pre-Mesozoic Geology in the Subsurface of Peel River Map Area, Yukon Territory and District of Mackenzie. Geological Survey of Canada, Memoir 401, 61 p.

Snowdon, L.R., Brooks, P.W., Williams, G.K. and Goodarzi, F., 1987. Correlation of the Canol Formation source rock with oil from Norman Wells. Organic Geochemistry, vol. 11, no. 6, p. 529-548.