PSApplying Regional Chemostratigraphic Variability to Refine Sequence Stratigraphic Frameworks within the Woodford Shale, Oklahoma*

Bryan W. Turner¹, Jessica A. Tréanton¹, and Roger M. Slatt¹

Search and Discovery Article #51040 (2014)**
Posted December 15, 2014

*Adapted from poster presentation given at the Geoscience Technology Workshop, Permian and Midland Basin New Technologies, Houston, Texas, September 4-5, 2014
**AAPG©2014 Serial rights given by author. For all other rights contact author directly.

¹University of Oklahoma, Norman, OK, USA (bwturner@ou.edu, jessica.treanton@ou.edu)

Abstract

The Woodford Shale provides an opportunity to test recent advances in handheld XRF (HHXRF) technology to develop sequence stratigraphic frameworks by comparing chemostratigraphic profiles directly to gamma ray logs and measured stratigraphic sections obtained from the same locations. Gamma ray profiles allow a direct integration of these newly obtained chemostratigraphic profiles into previously interpreted sequence stratigraphic frameworks. One field site and three cores of the Woodford Shale have been scanned using the HHXRF to develop a preliminary analysis of the regional variability of chemostratigraphic profiles for this formation.

Three cores from Caddo, Washington, and Pottawatomie Counties in Oklahoma and one outcrop at the Hunton Anticline Quarry (HAQ) in Murray County, OK represent both proximal and distal environments of the Woodford Shale. Clean surfaces at each area are scanned at one-foot intervals using the HHXRF to determine the elemental profiles. At the same resolution, a gamma ray profile is scanned using a GR scintillometer or core spectral gamma ray. The lithologic description, gamma ray profile, and elemental profiles are then used to develop the sequence stratigraphic interpretation.

Correlations based on regional scale variations in chemostratigraphic proxies show an overall increase in sedimentation in proximal settings with a decrease in sedimentation in distal settings. Drops in continentally derived sedimentation are interpreted from dilution of the Zr and Ti signals, associated with clastic sedimentation, relative to the concentration of K and Al, which are associated with clay mineral accumulation. All these elements are regarded as relatively immobile elements. In the distal setting, the chemostratigraphic ratios between clastic and clay proxies show a dilution of Ti and Zr between 10-60% in the upper Woodford relative to the lower Woodford, while in the proximal setting these ratios are concentrated up to 12-15% in the upper Woodford relative to the lower. These changes are indicative of a landward shift in sedimentation.

pyright Bryan Turner 2014. For reproduction rights contact author.

APPLYING REGIONAL CHEMOSTRATIGRAPHIC VARIABILITY TO REFINE SEQUENCE STRATIGRAPHIC FRAMEWORKS WITHIN THE WOODFORD SHALE, OKLAHOMA

TURNER, Bryan W.1*, TRÉANTON, Jessica A.1+, and SLATT, Roger M.2

(1) ConocoPhillips School of Geology and Geophysics, University of Oklahoma, 100 East Boyd St., Office 912, Norman OK 73019 (2) ConocoPhillips School of Geology and Geophysics, University of Oklahoma, 100 East Boyd St., Office 934, Norman OK 73019




AAPG Permian and Midland Basin New Technologies


Abstract:

The Woodford Shale provides an opportunity to test recent advances in handheld XRF (HHXRF) technology to develop sequence stratigraphic frameworks by comparing chemostratigraphic profiles directly to gamma ray logs and measured stratigraphic sections obtained from the same locations. Gamma ray profiles allow a direct integration of these newly obtained chemostratigraphic profiles into previously interpreted sequence stratigraphic frameworks. One field site and three cores of the Woodford Shale have been scanned using the HHXRF to develop a preliminary analysis of the regional variability of chemostratigraphic profiles for this formation.

Three cores from Caddo, Washington, and Pottawatomie Counties in Oklahoma and one outcrop at the Hunton Anticline Quarry (HAQ) in Murray County, OK represent both proximal and distal environments of the Woodford Shale. Clean surfaces at each area are scanned at one foot intervals using the HHXRF to determine the elemental profiles. At the same resolution, a gamma ray profile is scanned using a GR scintillometer or core spectral gamma ray. The lithologic description, gamma ray profile, and elemental profiles are then used to develop the sequence stratigraphic interpretation.

Correlations based on regional scale variations in chemostratigraphic proxies show an overall increase in sedimentation in proximal settings with a decrease in sedimentation in distal settings. Drops in continentally derived sedimentation are interpreted from dilution of the Zr and Ti signals, associated with clastic sedimentation, relative to the concentration of K and Al which are associated with clay mineral accumulation. All these elements are regarded as relatively immobile elements. In the distal setting the chemostratigraphic ratios between clastic and clay proxies show a dilution of Ti and Zr between 10-60% in the upper Woodford relative to the lower Woodford, while in the proximal setting these ratios are concentrated up to 12-15% in the upper Woodford relative to the lower. These changes are indicative of a







approximately 100 feet in the HAQ stratigraphic column, there is an anomalous spike is Si in relation to Al. This is indicative of SiO₂ deposition. As Ti and Zr are also low at this horizon, it is interpreted to be iogenic in origin, indicating a depositional hiatus.

500 1000

GR (cps)

spike in carbonate proxies (Ca and Sr). This could be analogous to reciprocal sedimentation (Schlager, 2005) where carbonate accumulation indicates clastic sedimentation is sequestered landward. A) A thin section showing the biogenic quartz. B) The hand sample showing the biogenic quartz horizon.





within continentally derived sediments and other elements are associated with clar minerals (Rodrigues, 2007). By comparing these elemental concentrations, it is approximate the redox state at the sediment water interface (Alaeo and Maynard. 204; Brumsack, 2006; Tribovillard et al., 2006). However, it is important to utilize multiple elemental proxies (Rowe et al., 2012), preferably of low mobility, to limit

understood to be a part of sequence stratigraphic frameworks (e.g. Haq et al.,

Finally, subtle surfaces of non-deposition have been interpretted to occur on horizons where carbonate proxies (Ca and Sr) peak or where the Si/Al rato peaks while other continental sedimentation indicators remain constant.

Conclusions:

Three cores from Caddo, Washington, and Pottawatomie Counties in Oklahomo and one outcrop at the Hunton Anticline Quarry (HAQ) in Murray County, OK represent both proximal and distal environments of the Woodford Shale. The develop the sequence stratigraphic interpretation. Regional correlations of primarily based on Gamma Ray profiles. Application of chemostratigraphy co reveal additional detail to help clarify correlations where ambiguity may be present (Tréanton et al., this conference).

Nested cyclicity has long been understood to be a component of sequence stratigraphy (e.g. Hag et al., 1987). The Gamma ray log from the core Hall 2B reveals that there is an overall cleaning upward succession. This can be interpreted as an overall progradation of sediments associated with a regression of the shorelline. However, chemostratiaraphy clearly shows that each of the shallowing upward cycles is composed of shorter periods of localized rises and falls in stratigraphic base level.

Chemostratigraphy can also reveal important details that may be absent from traditional well logs. The Anthis 2 well log shows a gamma ray log with very little with each succession capped by a spike in carbonate production. These successions appear to be analogous to reciprocal sedimentation (Schlager, 2005).

chemostratianaphy. Gamma ray correlation between the Ray 1-13 and the HAQ is relatively straightforward. However superimposing the chemostratigraphic cyclicity on the gamma ray correlation reveals that there are localized events nested within the regional trends.

shales of Upper Pennsylvanian Kansas-type cyclothems. Chem. Geo., 206,

Pennsylvanian Kansas-type cyclothems. Palaeogeog. Palaeoclim. Palaleoecol., 232, Haa, B. U., Hardenbol, J., and Vail, P.R. (1987) Chronology of Fluctuating Sea Levels

Since the Triassic. Science v. 235, p. 1156-1166. Rodrigues, R. (2007) Chemostratigraphy. In Topics in Geobiology vol. 23 Applied

Stratigraphy (ed. Koutsoukos, E.A.M.) Springer, Dordrecht. p 165-178. Rowe, H., Hughes, N., and Robinson, K. (2012) The quantification and application

chemostratigraphy and geochemistry. Chem. Geo., 324-325, p. 122-131. chalger, W. (2005) Sequence Stratigraphy of the T Factory in Carbonate

Sedimentology and Sequence Stratigraphy. SEPM Concepts in Sedimentology and

réanton, J., Turner, B., and Slatt, R. (2014) Outcrop Derived Inorganic Geochemistry of the Woodford Shale; Murray County, Oklahoma. Houston Geological Society Mudrocks Conference, Feb 17-18, Houston, TX.

We would like to thank Matt Pranter for allowing us to use his hand-held training on the Niton HH-XRF model, Brian Cardott for

