

# **Spatial and Temporal Deformation of a Lake Malawi Accommodation Zone: Integration of Dated Scientific Drill Cores and Seismic Data\***

**Tannis McCartney<sup>1</sup> and Christopher A. Scholz<sup>1</sup>**

Search and Discovery Article #51048 (2014)

Posted December 8, 2014

\*Adapted from extended abstract prepared in conjunction with presentation at CSPG/CSEG/CWLS GeoConvention 2013, (Integration: Geoscience engineering Partnership) Calgary TELUS Convention Centre & ERCB Core Research Centre, Calgary, AB, Canada, 6-12 May 2013, AAPG/CSPG©2014

<sup>1</sup>Department of Earth Sciences, Syracuse University, Syracuse, NY ([tmccartn@syr.edu](mailto:tmccartn@syr.edu))

## **Abstract**

The Lake Malawi Rift is located in the southern extension of the western arm of the East Africa Rift System ([Figure 1a](#)). At its northern end, this system contains the Albertine Graben ([Figure 1a](#)), where a large hydrocarbon province was recently discovered.

## **Method**

Three vintages of seismic data have been collected in Lake Malawi. Project PROBE (1980s) collected low-resolution, high-penetration multi-channel seismic (six second records). The Central Basin was the focus of a series of nested single-channel seismic surveys in the 1990s. In 2001, 4-second multi-channel seismic with moderate resolution was acquired. The single-channel data, collected using a small air gun with a bandwidth of 50-500Hz, is higher resolution and has greater fidelity than the multi-channel seismic acquired in the same region. An integrated interpretation of the single-channel seismic and the Project PROBE seismic is presented here.

## **Discussion**

Structural domains can be identified both laterally and temporally in the migrated single-channel seismic. On the hanging wall of the Central Basin border fault is a segmented accommodation zone characterized by a network of shallow synthetic and antithetic faults. This accommodation zone was uplifted in the last 50,000 years. It is bounded to the northwest by a single fault and to the southeast by several linear fault segments. A rotated block beneath these faults is evidence for faulting that occurred over one million years ago. The flanks of the accommodation zone contain a basin-fill facies (Lyons et al., 2011). Deformation on the northwest flank is associated with mud diapirs that have appeared in the last 50,000 years. The southeast flank contains several faults but no apparent diapirs.

## **Conclusions**

The faults in the accommodation zone have a complex geometry; understanding their evolution is key to understanding the inception, subsidence, and propagation of continental rifts. New information from an actively deforming accommodation zone will help constrain an important part of rift basin petroleum systems.

## **References Cited**

Amante, C., and B. Eakins, 2009, ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis: NOAA Technical Memorandum NESDIS NGDC-24, 19 p.

Lyons, R.P., C. Scholz, M.R. Buoniconti, and M.R. Martin, 2011, Late Quaternary stratigraphic analysis of the Lake Malawi Rift, East Africa: An integration of drill-core and seismic-reflection data: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 303/1-4, p. 20–37, doi: 10.1016/j.palaeo.2009.04.014.




Figure 1. ETOPO1 1 Arc-Minute global relief maps (Amante and Eakins, 2009). (A) The western branch of the East African Rift System. Lake Albert, at the north end of the branch, is in the Albertine Graben. Lake Malawi is in the south. (B) Lake Malawi, with the three principle half-graben basins circled (border faults in blue).