

PS Short Junction Field Core Description and Petrophysical Analysis of the Hunton Group, Cleveland County, Oklahoma*

Tim Hunt³, John Speight¹, Valentina Vallega², Huabo Liu¹, Julio Garcia², and Curtis Helms⁴

Search and Discovery Article #20371 (2016)**

Posted October 17, 2016

*Adapted from poster presentation given at AAPG Southwest Section Meeting, Abilene, Texas, April 9-12, 2016

**Datapages © 2016 Serial rights given by author. For all other rights contact author directly.

¹Trey Resources, Inc. (thuntgeo@gmail.com)

²Schlumberger

³retired

⁴Midland College, PPDC

Abstract

Two units comprise most of the Short Junction Field, which produces from the Hunton Group, located in northeast Cleveland County, Oklahoma. The units have produced approximately 22 million barrels of oil since 1948 of an estimated 250 million OOIP. The less than 9% recovery even after a secondary water flood leaves a sizable target for a revitalized field.

Trey Resources, Inc. took over operations in 2014 and drilled the WSJU 1101H. The entire Hunton (Bois d'Arc to Chimneyhill) was cored, as well as a full petrophysical suite including borehole imaging logs. In 2008, the WSJU 109H, was recompleted as a horizontal lateral and included borehole imaging logs. These data were used to model the Bois d'Arc.

The core was oriented to determine principle stress direction and structural position. Additional whole core samples were analyzed for directional permeability and plugs samples were measured permeability in the east-west direction. Three plugs were selected for conventional CT scan analysis to help determine electrical properties.

Advanced interpretation techniques were applied on the acquired borehole images and correlated with the core results. The objective was to characterize the heterogeneities present in the formation. With the creation of full borehole images covering the entire borehole surface, it was possible to better identify various heterogeneities (including vugs and fractures) and classify them as connected or isolated vugs, fractures connecting vugs, or heterogeneity developed along bed boundaries. Intervals where the matrix porosity was the predominant component to the overall porosity were highlighted, versus intervals where the vuggy porosity has an important contribution.



Short Junction Field Core Description and Petrophysical Analysis of the Hunton Group, Cleveland County, Oklahoma

Tim Hunt³, John Speight¹, Valentina Vallega², Huabo Liu¹, Julio Garca², Curtis Helms⁴ (1) Trey Resources, Inc., (2) Schlumberger, (3) Retired, (4) Midland College, PPDC




Abstract

Two units comprise most of the Short Junction field, which produces from the Hunton group, located in northwest Cleveland County, Oklahoma. The units have produced approximately 22 million barrels of oil since 1948 of an estimated 250 million OOIP. The less than 9% recovery even after a secondary water flood leaves a sizable target for a revitalized field.


In 2008, the WSJU 109H, was recompleted as a horizontal lateral and included borehole imaging logs. Trey Resources acquired the units in 2014 and drilled the WSJU 1101H. The entire Hunton (Bois d'Arc to Chimneyhill) was cored as well as a full petrophysical suite including borehole imaging logs. These data were used to model the Bois d'Arc.

The core was oriented to determine principle stress direction and structural position. Additional whole core samples were analyzed for directional permeability and plugs samples were measured permeability in the east west direction. Three plugs were selected for conventional CT scan analysis to help determine electrical properties.

Advanced interpretation techniques were applied on the acquired borehole images and correlated with the core results. The objective was to characterize the heterogeneities present in the formation. With the creation of full borehole images covering the entire borehole surface, it was possible to better identify various heterogeneities (including vugs and fractures) and classify them as connected or isolated vugs, fractures connecting vugs or heterogeneity developed along bed boundaries. Intervals where the matrix porosity was the predominant component to the overall porosity were highlighted, versus intervals where the vuggy porosity has an important contribution.




Short Junction Location Map




Simplified Stratigraphic Column				
Short Junction Field				
System	Series	Stage	Group	Formation
Permian	Leonardian			Garber - Wellington
	Wolfcampian			Neva
Pennsylvanian	Virgilian	Shawnee		Pawhuska
				Hoover
				Tonkawa
	Missouri	Skiatook		Hogshooter
				Checkerboard
	Des Moines	Marmaton		Big Lime
				Oswego
		Cherokee		Prue
				Verdigris
				Skinner
				Pink Lime
				Red Fork
				Inola
				Bartlesville
			Atoka	
	Morrow			
Mississippian	Chester			Absent @ Short Junction
	Meramec			
	Osage			
	Kinderhook			
Devonian				Woodford
	Silurian	Hunton Grp		Misener
				Frisco
			Bois d'Arc/Haragan	
			Henryhouse	
Ordovician	Chimneyhill Subgroup		Clarita	
			Cochrane	
			Keel	
	Simpson Grp		Sylvan	
			Viola	
			Bromide/Wilcox	
			Tulip Creek	
			McLish	
			Oil Creek	
			Joins	
Cambrian	Atoka Grp			
			Reagan	
Precambrian				Granitic Basement


Trey Resources, Inc West Short Junction Unit #1101H



Core description WSJU #1101 Page 1 of 4



Core description WSJU #1101 Page 2 of 4





Short Junction Field Core Description and Petrophysical Analysis of the Hunton Group, Cleveland County, Oklahoma

Tim Hunt³, John Speight¹, Valentina Vallega², Huabo Liu¹, Julio Garca², Curtis Helms⁴ (1) Trey Resources, Inc., (2) Schlumberger, (3) Retired, (4) Midland College, PPDC

Core description WSJU #1101 Page 3 of 4

Core description WSJU #1101 Page 4 of 4

WSJU #1101

FMI Interpretation

On the left, part of the PorosTox results, applied to the WSJU 1101H

- Track 1: Depth Reference, scale 1:24
- Track 2: Zonations highlighting intervals used for setting thresholds
- Track 3: Full Calibrated image
- Track 4: Calibrated FMI² Microseismicity curve with 0.2' vertical resolution
- Track 5: Heterogeneity image delineation. Refer to legend top of log.
- Track 6: Image Porosity Map. Increase in darkness equals increase in image porosity
- Track 7: Spectrum of porosity distribution
- Track 8: Cumulative porosity distribution
- Track 9: Average image porosity at each heterogeneity type
- Track 10: Cross plot porosity and total porosity computed from image porosity.

Value corresponds to average at each depth level of image porosity curve.

Image Porosity Analysis Workflow

This state of the art workflow includes textural analysis, image porosity analysis and fracture analysis to fully characterize the porosity distribution in the carbonate reservoir. This technology was applied in the West Short Junction Unit 1101 and in the West Short Junction Unit 109H.

- 1) Full image creation: this step utilizes geostatistics to generate an image that represents full borehole coverage
- 2) Conductive and Resistive heterogeneities are delineated utilizing thresholds on contrast and resistivity values. Changes in resistivities compared to the matrix corresponds to heterogeneities: highly resistive heterogeneities correspond to cemented zones, while low values of resistivities correspond to vugs or fractures (Delhomme, 1992)
- 3) Combining the detailed features identification done in the manual dip picking phase, with the heterogeneity delineation, allow the classification of heterogeneities in different categories.
- 4) Porosity map from image is constructed utilizing a well established method which computes porosity from a modified Archie's equation applicable to the flushed zone and having as input each conductivity curve's measure by the Formation Micro Imager (Newberry et al, 1996)







Image to core comparison

core description
fault fillings, and
karst related features

Core description
karst filling breccias




Response of Porotex in Deep Shelfal environment



The general response in Deep shelfal environment is lack of heterogeneities and overall porosity from image from low to medium

Response of Porotex in Organic Mud Mound



The general response in the organic mud mound is of presence of heterogeneities contributing to the overall medium image porosity. Heterogeneities are predominantly along beddings and isolated

Response of Porotex in Shallow Shelf




The general response in shallow shelf environment is abundance of heterogeneities along beddings and overall porosity from image low



Short Junction Field Core Description and Petrophysical Analysis of the Hunton Group, Cleveland County, Oklahoma

Tim Hunt³, John Speight¹, Valentina Vallega², Huabo Liu¹, Julio Garca², Curtis Helms⁴ (1) Trey Resources, Inc., (2) Schlumberger, (3) Retired, (4) Midland College, PPDC

Formation Micro Imager of horizontal lateral West Short Junction Unit #109H




Overall a total of 5 FMI image facies were identified and below is represented the output of the porosity classification analysis:


- 1) High background matrix resistivity with discrete fracture presence
- 2) Medium to High background resistivity with high presence of fractures
- 3) Medium background resistivity with segmented fractures
- 4) Low background resistivity with vuggy texture
- 5) Low background resistivity with vuggy texture and fractures




FACIES 1
Low presence of heterogeneities
Discrete fractures are the contributors to reservoir properties




FACIES 2
Low presence of heterogeneities and if present connected to fractures
High presence of fractures is the contributor to reservoir properties



FACIES 3
Increased presence of heterogeneities of various nature
Segmented fractures and heterogeneities equally present



FACIES 4
Increased presence of heterogeneities and overall increased matrix porosity. Heterogeneities seem to be predominant along boundaries, connected or isolated.
Fractures do not represent a predominant feature



FACIES 5
Increased matrix porosity and increased heterogeneity presence. Vugs connected to fractures are the most predominant feature.
Fractures and vugs are equally highly contributing to increased reservoir properties