

Re-evaluation of the Arikareean (Late Oligocene to Early Miocene) Tecuya Canyon Land Mammal Local Faunas, Lower Member, Continental Tecuya Formation, Northern San Emigdio Mountains, Kern County, California

E. Bruce Lander

Paleo Environmental Associates, Inc., Altadena, CA

paleo@earthlink.net

The Tecuya Canyon Local Faunas (TCLFs) consist of two stratigraphically superposed fossil land mammal assemblages that occur in the lower mbr. of the continental Tecuya Fm. The fossil-bearing strata grade laterally into the marine upper Pleito and stratigraphically overlying lower Temblor Fms., which contain fossil marine molluscan and benthic foraminiferal assemblages.

The Lower TCLF occurs roughly 75 m below the Tunis Volcanics at CIT locality 116 and UCMP localities 3352 and 3354; includes *Archaeolagus* sp. (rabbit), *Protosciurus tecuyensis* (Bryant) (squirrel), *Leidymys nematodon* (Cope) (mouse), *Daphoenus* sp. (includes *Daphoenus? ruber* Stock) (bear dog), *Miohippus annectens* Marsh (horse), *Diceratherium armatum* Marsh (large rhinoceros), *Eporeodon major* (Leidy) (mediumsized oreodont), *Nanotragulus loomisi* Lull (small brachyodont mouse deer), and *Hypertragulus planiceps* (Sinclair) (large mouse deer); and is from a stratigraphic interval probably laterally equivalent to one containing mollusks assignable to an unnamed stage and lower Zemorrian benthic foraminifers. The overlapping biochronologic ranges of *Archaeolagus*, *L. nematodon*, *D. annectens*, *E. major*, *N. loomisi*, and *H. Planiceps* indicate the Lower TCLF is late early Arikareean in age and a correlative of the Blakeley, Alamos Canyon, and upper South Mountain Local Faunas (LFs) (uppermost upper mbr. Sespe Fm. [SF]) in Ventura Co., the Lower Bowerman Landfill LF (lowermost upper mbr., SF) in Orange Co. (OC), and the Upper Eastlake LF (sandstone-mudstone mbr., Otay Fm.) in San Diego Co.

The Upper TCLF occurs only about 8-9 m below the volcanics at UCMP locality V-6378; includes *Oreolagus* sp. (pika) and *L. nematodon*, which have overlapping biochronologic ranges only during the late early and late Arikareean; and possibly is a correlative of the late Arikareean Upper OC Sespe/Vaqueros Fauna (lower [but not lowermost] upper mbr., SF). The Upper TC LF is from a stratigraphic interval laterally equivalent to one containing mollusks assignable to the Vaqueros Stage and upper Zemorrian benthic foraminifers.