

## **The Early Jurassic Heracles Sequence, Scotian Basin, Canada: Recognition of Latest Stage Synrift Tectonism and Correspondence to Structures Offshore Morocco**

**David E. Brown<sup>1</sup>, Sonya A. Dehler<sup>2</sup>, Keith Loudon<sup>3</sup>, and Yue Wu<sup>4</sup>**

<sup>1</sup> Canada-Nova Scotia Offshore Petroleum Board, 1791 Barrington Street, Halifax, Nova Scotia, B3J 3K9, Canada

<sup>2</sup> Geological Survey of Canada (Atlantic), Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada  
sdehler@nrcan.gc.ca

<sup>4</sup> Department of Oceanography, Dalhousie University, Halifax, Nova Scotia, B3H 4J1, Canada

Seismic profiles on the Scotian shelf, slope and abyssal plain offshore Eastern Canada reveal a previously unrecognized earliest Jurassic post-salt / pre-breakup stratigraphic succession. The Heracles Sequence is observed on the shelf margin as an eastward-directed infill succession within a series of half grabens having counter-regional, northwest-dipping boundary faults. On the slope, its inferred presence in the salt depocentre adjacent to the basin hingeline is masked by a thick wedge of later Mesozoic and Cenozoic sediments and salt structures. In deep water, it is recognized as a poorly-imaged but apparently extensive sequence between interpreted basement and the late Sinemurian breakup unconformity. At the eastern extremity of this region, it appears as westward-thickening wedges in highly rotated fault blocks.

The Heracles Sequence is interpreted as the product of the last phase of synrift tectonism prior to separation of the Nova Scotian and Moroccan conjugate margins in the late Sinemurian. Post-salt (early Hettangian) uplift of the mainland Nova Scotia shoulder region and the eventual rift spreading centre provided sources for sediments that prograded east- and westwards respectively into the main salt basin. Interpreted fluvial sequences advanced over marine evaporates ponded in depressions on a rifted basement setting and induced syndepositional halokinesis and the formation of salt-evacuation synclines. Where thin on the margins (especially near the future spreading axis), the salt provided a detachment surface and facilitated the observed high rotation on loading-induced fault blocks during a final uplift phase. This interpretation buttresses other geophysical evidence that suggests the underlying basement may not be oceanic crust, as previously proposed, but rather highly attenuated and fractured continental crust or serpentinitized mantle.

The recognition of this late stage pre-breakup synrift sequence in the Scotian Basin offers important insights on this phase of the rifting process, and possibly its Moroccan conjugate and other margins. It thus has significant implications regarding the recognition of the continental crust and crustal boundaries, age and timing of syntectonic deposition and salt tectonism, original distribution and extent of marine evaporite sequences, timing and style of rifting, modeling of crustal heat flow, and petroleum systems attributes and modeling.