
Sequence Stratigraphic Framework and Depositional History of Oligocene Frio Slope-Fan, Lowstand Prograding Wedge, and Shallow-Marine Transgressive-Regressive Deposits in the Lavaca Bay Area, Texas

William A. Ambrose¹, Brandon Johnson¹, Ursula Hammes¹, and Dean Johnstone²

¹Bureau of Economic Geology, The University of Texas at Austin,
University Station, Box X, Austin, Texas 78713-8924

²Neumin Production Company, P.O. Box 769, Point Comfort, Texas 77978

ABSTRACT

The Frio Formation in Lavaca Bay, Texas, provides an excellent example of a stratigraphic succession of third-order lowstand slope fans overlain by lowstand progradational wedges capped by high-frequency transgressive-regressive cycles. Frio reservoir compartments are controlled by narrow sandstone-body geometry, wedging, and pinchouts of sandstones on the crest of rollover structures, deep shale mounding, and structural complexity from syndepositional growth faults and closely-spaced antithetic faults. The lower Frio Formation contains three 800- to 1,000-ft (240- to 330-m), third-order slope-fan sequences composed of amalgamated channel and lobate fan deposits. Slope-channel axes are commonly deflected along the strike of growth faults, suggesting syndepositional fault movement that locally modified slope bathymetry. The upper and middle Frio Formation contains a lowstand prograding-wedge sequence with low-angle (<5°) clinoforms that downlap onto a prominent angular unconformity (29.0 Ma) at the top of geopressure. In contrast, the upper Frio Formation contains high-frequency, shallow-marine fourth- and fifth-order transgressive-regressive cycles in an overall third-order transgressive systems tract. These transgressive-regressive cycles exhibit continuous sandstone-body geometries, are structurally simple, and therefore have less potential than the lower and upper Frio Formation for reservoir compartmentalization.

Ambrose, W. A., B. Johnson, U. Hammes, and D. Johnstone, 2010, Sequence stratigraphic framework and depositional history of Oligocene Frio slope-fan, lowstand prograding wedge, and shallow-marine transgressive-regressive deposits in the Lavaca Bay area, Texas: Gulf Coast Association of Geological Societies Transactions, v. 60, p. 29-38.