

Geodynamic evolution of Tethys in the Lesser Caucasus: deciphering clues from Radiolarian biochronology

Danelian T.¹, Asatryan G.^{2,3}, Galoyan G.³, Sahakyan L.³, and Sosson M.⁴

¹ University Lille 1, Department of Earth Sciences, CNRS-FRE 3298 "Géosystèmes", Bât SN5, 59655 Villeneuve d'Ascq, France (<mailto:taniel.danelian@univ-lille1.fr>)

² Institut of Geological Sciences, Armenian National Academy of Sciences, 24a Baghramian avenue, Yerevan, 375019, Armenia,

³ University Pierre & Marie Curie (Paris VI), CNRS-UMR 7207 Centre de Recherches sur la Paléobiodiversité et les Paléoenvironnements (CR2P), C. 104, 4 place Jussieu. 75005 Paris, France

⁴ University of Nice – Sophia Antipolis, UMR Géoazur, CNRS, OCA, 250 rue A. Einstein. 06560 Valbonne 2, France

Radiolaria are often the only fossils preserved in the sedimentary cover of Tethyan ophiolites. They have thus become an important tool for the investigation of the geodynamic evolution of Alpine mountain belts.

The main objective of our project is to specify the geohistory of ophiolites preserved in the Lesser Caucasus, a key area of the Alpine-Himalayan mountain belt. Improved knowledge of the age of Caucasian ophiolites will allow better lateral correlations of possibly equivalent suture zones in Turkey and Iran and will help deciphering the geodynamic evolution of the greater area between Eurasia and the South-Armenian Block. The latter was a micro-continent that has become detached of Gondwana during the Late Palaeozoic – Early Mesozoic. It is considered as part of the eastward extension of Taurides-Anatolides continental microplate.

Two main ophiolitic zones are recognised in the Lesser Caucasus:

- The Amassia-Sevan-Akera ophiolites (running from NW Armenia, through East of Lake Sevan to western Karabakh) represent the main suture zone in the Lesser Caucasus, including extensive outcrops of peridotites (often serpentinitized) and gabbros. The geochemistry of lavas suggests the presence of two distinct extrusive systems; i) a contaminated MORB series of basalts, bearing a slight calc-alkaline signature, is considered to be the result of a slow spreading ridge in a back-arc setting; ii) an alkaline OIB series of lavas considered to be the expression of a mantle plume event.
- The Vedi ophiolite, in the SE of Yerevan, is also composed of serpentinites, gabbros and a thick pile of massive and pillowed lava flows. It is considered as a folded klippe sequence that was thrust above Cenomanian-Turonian shallow water carbonates and flysch of the South Armenian Block.

New and revised radiolarian data point to the following working scenario:

- (1) the initial phase of sea floor spreading took place during the Late Triassic (Carnian). Evidence for this comes from a single locality (Zod pass, Knipper et al. 1997);
- (2) The bulk of oceanic crust, preserved today in both the Sevan-Akera and Vedi ophiolite complexes, was formed during the Middle Jurassic (Bajocian, Bathonian-Callovian) in a supra-subduction zone (SSZ) setting;
- (3) Submarine volcanic activity continued occasionally during the Late Jurassic and Early Cretaceous, as this is suggested by reliable radiolarian ages obtained on cherts intercalated with tholeiitic lavas from both the Vedi and Sevan ophiolites.
- (4) Preliminary results on radiolarian cherts intercalated with alkaline lavas of the Sevan ophiolite suggest that a hot spot event took place during the Late Jurassic.
- (5) Several tuffite levels, intercalated within the radiolarite sequence of the Sevan ophiolite, reflect a Late Jurassic subaerial volcanic. At the moment, it is unclear whether they are related with the emplacement of the SSZ ophiolites or with the mantle plume event.