

Episodic Tectonics in the Phanerozoic Succession of the North American Arctic and the “10 Million Year Flood”

Ashton Embry, Geological Survey of Canada, Calgary, aembry@nrcan.gc.ca

Benoit Beauchamp, University of Calgary, Calgary, bbeauch@ucalgary.ca

Keith Dewing, Geological Survey of Canada, Calgary, kdewing@nrcan.gc.ca

James Dixon, Geological Survey of Canada, Calgary, jdixon@nrcan.gc.ca

Introduction

The goal of this study is to catalogue all the large scale, tectonically-generated sequence boundaries which punctuate the Phanerozoic stratigraphy of the North American Arctic and to determine the frequency and possible origin of the interpreted tectonic episodes which gave rise to the boundaries. This study is feasible because, over the last 40 years, the Phanerozoic succession of the North American Arctic, both on the surface and in the subsurface, has been intensely studied in various sedimentary basins with modern basin analysis techniques including sedimentology, sequence stratigraphy and biostratigraphy.

Data Sources

Notably, the entire Phanerozoic succession is present in the NA Arctic although no single basin contains a complete column. The data for this study come mainly from three major basins: Franklinian Basin, Sverdrup Basin and Beaufort-Mackenzie Basin. Data from other basins in the NA Arctic such as the North Slope Basin of Alaska have been used to complement and refine the results obtained from the three main basins.

The Franklinian Basin of the Canadian Arctic Islands and adjacent northern Greenland provides an excellent record of the Cambrian to Devonian succession from basin margin to basin centre and the stratigraphy of the succession is detailed in numerous reports, including Mayr (1978), Thorsteinsson and Mayr (1987), Higgins et al. (1991), Trettin et al. (1991), Embry (1991a), Mayr et al (1994), Harrison (1995), Mayr et al. (1998), de Freitas et al. (1999), and Dewing et al. (2008). Relevant sequence stratigraphic data on the Cambrian-Devonian succession have also been gleaned from work done on the northern Canadian mainland where the strata are widely distributed (Fritz et al., 1992; Pyle and Jones, 2009).

Stratigraphic data for the Carboniferous through Cretaceous succession have been obtained primarily from the Sverdrup Basin of the Canadian Arctic Archipelago. The succession is very thick (12 km) and exquisitely exposed in the eastern part of the basin. In the west, ample well and seismic data supplement the surface exposures. The key references for the detailed stratigraphy of this succession are Balkwill (1978, 1983), Beauchamp and Henderson (1994), Beauchamp and Thériault (1994), Beauchamp (1995), Beauchamp et al. (2001, 2009), Beauchamp and Olchowy (2003), Embry (1991b, 1993, 1997, 2011) and Embry and Beauchamp (2008). Data for the Jurassic and Cretaceous succession were also acquired from studies in the Mackenzie Delta and adjacent areas (Dixon 1982, 1991, 1993; Poulton, 1982), offshore west Greenland (Schenk, 2011) and the North Slope Basin of Alaska (Houseknecht and Bird, 2004; Decker, 2007; Bird and Houseknecht, 2011).

Cenozoic stratigraphic data are most complete for the Beaufort–Mackenzie Basin on the northern continental margin. The succession is mainly in the subsurface and has been unraveled through the analysis of plentiful well and seismic information. Important references for the Cenozoic stratigraphy include Dixon et al (1992), Dixon (1996), Dixon et al. (2008) and Helwig et al (2011). Additional stratigraphic data were obtained from the outcropping Paleogene succession of the eastern Sverdrup Basin (Ricketts, 1991, 1994; Ricketts and Stephenson, 1994; Harrison et al., 1999).

Tectonic Episodes

These detailed and basin-wide stratigraphic studies in the various basins have revealed the presence of 56, large magnitude, sequence boundaries within the Phanerozoic succession of Arctic North America. These boundaries are distinguished by the presence of the following characteristics:

- 1) A widespread, often angular, unconformity on the basin margins and positive elements
- 2) A major change in depositional regime
- 3) A notable change in tectonic regime and subsidence pattern
- 4) A change in source region for siliciclastic sediments
- 5) A major back step of the carbonate platform edge for carbonate sediments
- 6) A widespread transgression with significant deepening directly following the boundary

The sequence boundary characteristics strongly indicate that the boundaries identified were generated primarily by tectonics and not by eustasy (Embry, 1990). Thus, each sequence boundary is interpreted to be generated by a tectonic episode which affected the entire region and which resulted in tectonic uplift and erosion on the margins of the basins. This uplift was accompanied by notable regression in the more central areas of the basins and was followed by a collapse and marine flooding of the basin margins and beyond. The sequence boundaries, which punctuate the Phanerozoic stratigraphy of the NA Arctic, occurred on average about every 10 million years (56 boundaries in 545 million years).

The maximum regressive surface portion of each sequence boundary has been dated on the basis of the available biostratigraphy which, in most cases, allows the boundary to be assigned to a specific biozone. The sequence boundaries were then assigned a numerical age through the correlation of the biozones with the 2012 geological time scale (Gradstein et al., 2012). The time intervals between the boundaries were no less than 5 and no more than 15 million years in almost all cases. It is clear that the tectonically generated boundaries were not strictly periodic but were chaotic and had a relatively high chance of reoccurring within 10 million years. Furthermore, there is no systematic change in boundary frequency throughout the entire Phanerozoic and an average frequency of about 10 million years occurs for any given 100 million year interval. Thus, the major transgression which follows each boundary could perhaps be referred to as a “10 million year flood”.

A given tectonic episode began with the initial uplift of the basin margin (start of base level fall) and ended with the collapse and marine flooding of the margin (maximum flooding surface). The sequence boundary was generated during the tectonic episode and represents the time of maximum uplift and basinward extent of the unconformity. It is estimated that the duration of each of the tectonic episodes was in the range of a few million years and was significantly shorter than the intervening times of tectonic quiescence.

Origin of Tectonic Episodes

The recognition of these tectonic episodes leads to questions such as “what is the primary cause of such repetitive episodes?” and “are they global in extent or just a characteristic of the North American Arctic?” It is postulated that such tectonic episodes are an expression of relatively rapid and substantial changes in the horizontal and vertical stress fields of the NA Arctic. Such stress changes would be possibly due to somewhat abrupt changes in the speed and/or direction of plate movements which

affected the NA Arctic. Given there is a complex feedback between plate distribution and dynamics with mantle convection and heat release (Rolf et al., 2012), the recognized tectonic episodes would presumably reflect perturbations in mantle convection intermittently occurring every 5 -15 million years due to a gradual buildup and consequent, episodic release of stress and heat.

In terms of the possible global extent of the recognized events, it is worth noting that all of the 18 major, tectonic, sequence boundaries recognized by Sloss (1988) for the Phanerozoic of the North American continent are present in our compilation for the Arctic. Furthermore, a cursory literature survey has indicated most of the tectonic sequence boundaries we have recognized are present in basins on different continents. However, in many cases, the published data are not sufficient to document a tectonic origin for a given boundary in a given basin. A more detailed literature search for the Triassic (Embry, 1997) has revealed that all the Triassic, tectonically-generated sequence boundaries recognized in the NA Arctic are present in numerous basins in North America, Europe and Asia and were also tectonically generated in those areas.

It has also been determined that 16 of the 18 Cambrian-Pliocene mass extinctions identified by Bambach (2006) coincide with an identified tectonic episode and that many of the tectonic episodes correlate with significant carbon isotope excursions (Saltzman and Thomas, 2012). Such correlations would imply that the episodes were the expression of a global phenomenon rather than one confined to the NA Arctic. Thus, it is quite possible that most, if not all, of the recognized, tectonically-generated, sequence boundaries of the NA Arctic represent episodic tectonics which occurred on a global scale. Given that all the tectonic plates are linked and a mantle-driven, major adjustment in the speed or direction of one plate would require compensatory movement changes in all the other plates, a phenomenon of episodic global tectonics occurring with a frequency of 5 -15 million years is a theoretical possibility.

Implications for Petroleum Exploration

These results have implications for petroleum exploration in that petroleum traps would have been formed, altered and sometimes breached during the tectonic episodes. Also, the movement of subsurface fluids would have been greatly influenced by such intermittent convulsions of the sedimentary column. Finally the recognition and dating of these tectonically-generated sequence boundaries in combination with their potential global distribution allows their occurrence to be predicted in unexplored sedimentary basins.

Conclusions

Fifty-six, large magnitude sequence boundaries have been delineated in the Phanerozoic succession of Arctic North America. The characteristics of the boundaries indicate that they were primarily generated by tectonics. The boundaries occur with an approximate 10 million year frequency (9.8 +/- 3.1). Each boundary was generated during a tectonic episode interpreted to reflect a mantle-driven, plate tectonic reorganization and consequent changes in regional stress fields. Such episodes likely lasted for a few million years and were separated by longer intervals of relative tectonic quiescence. There are indications that the recognized tectonic episodes affected basins throughout the world.

References

- Balkwill, H.R. 1978. Evolution of Sverdrup Basin. AAPG Bulletin, **62**, 1004- 1028.
- Balkwill, H.R. 1983. Geology of Amund Ringnes, Cornwall, and Haig-Thomas islands, District of Franklin. Geological Survey of Canada, Memoir 390, 76 p.
- Bambach, R. K. 2006. Phanerozoic biodiversity mass extinctions. Annual Review of Earth and Planetary Science, **34**, 127–155.

Beauchamp, B., 1995. Permian history of Arctic North America. *In* The Permian of Northern Pangea, Volume 2: Sedimentary Basins and Economic Resources, Scholle, P.A., Peryt, T.M., Ulmer-Scholle, D. S. eds., Springer-Verlag, Berlin, Heidelberg, 3–22.

Beauchamp, B., Harrison, J.C., Utting, J., Brent, T.A., Pinard, S., 2001. Carboniferous and Permian subsurface stratigraphy, Prince Patrick Island, Northwest Territories, Canadian Arctic. Geological Survey of Canada Bulletin 565, 93 pp.

Beauchamp, B., Henderson, C.M., 1994. The Lower Permian Raanes, Great Bear Cape and Trappers Cove formations, Sverdrup Basin, Canadian Arctic: stratigraphy and conodont zonation. *Bulletin of Canadian Petroleum Geology* 42, 562–597.

Beauchamp, B., Henderson, C.M., Grasby, S.E., Gates, L.T., Beatty, T.W., Utting, J., James, N.P., 2009. Late Permian sedimentation in the Sverdrup Basin, Canadian Arctic: The Lindström and Black Stripe formations, *Bulletin of Canadian Petroleum Geology*. V. 57, p. 167–191

Beauchamp, B., and Olchowy, B., 2003. Early Permian buildups (Tolkien reefs) associated with subaqueous evaporites, Canadian Arctic: a record of syn-tectonic to post-tectonic reciprocal uplift and subsidence. *In* Permo-Carboniferous Carbonate Platforms and Reefs, W.M. Ahr, P.M. Harris, W.A., Morgan and I.D. Somerville, eds., SEPM Special Publication No. 78 and AAPG Memoir 83, 133–153.

Beauchamp, B., Thériault, P., 1994. Late Paleozoic syn- and post-rift sequences on Grinnell Peninsula, Canadian Arctic (Sverdrup Basin): evidence for basin margin tectonic disturbances associated with sequence boundaries. *In* Pangea: Global Environments and Resources, Embry, A.F., Beauchamp, B., Glass, D.J. eds., Canadian Society of Petroleum Geologists Memoir 17, 199–217.

Bird, K.J., and Houseknecht, D.W., 2011, Geology and petroleum potential of Arctic Alaska, *In*: Spencer, A., Embry, A., Gautier, D., Sørensen, K., and Stoupakova, A., eds., Arctic petroleum geology: Geological Society of London Memoir 35, p. 485–499.

Decker, P. 2009, Brookian sequence stratigraphic correlations, Umiat to Milne Point Field, west-central North Slope, Alaska. State of Alaska, Preliminary Interpretive Report 2007-2, 21 p.

De Freitas et al., 1999 Silurian System of the Canadian Arctic Archipelago, BCPG, V. 47 p. 136-193

Dewing, K., U. Mayr, J. C. Harrison, and T. De Freitas, 2008, Upper Neoproterozoic to Lower Devonian stratigraphy of northeast Ellesmere Island, *in* U. Mayr, ed., The geology of northeast Ellesmere Island adjacent to Kane Basin and Kennedy Channel, Nunavut: Geological Survey of Canada Bulletin 592, p. 31–108.

Dixon, J. 1982: Jurassic and Lower Cretaceous subsurface stratigraphy of the Mackenzie Delta-Tuktoyaktuk Peninsula, N.W.T.; Geological Survey of Canada, Bulletin 349, 52 p.

Dixon, J. 1991: The Neocomian Parsons Group, northern Yukon and adjacent Northwest Territories; Geological Survey of Canada, Bulletin 406, 54 p.

Dixon, J., 1993, Regional unconformities in the Cretaceous of northwest Canada. *Cretaceous Research*, v. 14, pp. 17–38.

Dixon, J. (Editor), 1996, Geological Atlas of the Beaufort-Mackenzie Area: Geological Survey of Canada Miscellaneous Report 59, 173 p.

Dixon, J., Dietrich, J. R., and McNeil, D. H., 1992, Upper Cretaceous to Pleistocene Sequence Stratigraphy of the Beaufort-Mackenzie and Banks Island areas, northwest Canada :Geological Survey of Canada Bulletin 407, 90p.

Dixon, J., Dietrich, J., Lane, L. and McNeil, D., 2008, Geology of the Late Cretaceous to Cenozoic Beaufort-Mackenzie Basin, Canada. *In*: Miall, A. (ed.) Sedimentary Basins of the World, Vol. 5; The Sedimentary Basins of the United States and Canada. Elsevier, Amsterdam, 551-569.

Embry, A. 1990. A tectonic origin for third-order depositional sequences in extensional basins implications for basin modelling. *In*: Quantitative Dynamic Stratigraphy. T. Cross (ed.). Prentice Hall, p. 491-502.

Embry, A.F. 1991a. Middle-Upper Devonian clastic wedge of the Arctic Islands. *In*: H.P. Trettin (ed.), Geology of the Inuitian Orogen and Arctic Platform, Geological Survey of Canada, Geology of Canada, no. 3, p. 263-279.

Embry, A.F., 1991b. Mesozoic history of the Arctic Islands. *In*: H. Trettin (ed) Inuitian Orogen and Arctic Platform: Canada and Greenland. Geological Survey of Canada, Geology of Canada, 3, 369-433.

Embry, A.F., 1993. Transgressive-regressive (T-R) sequence analysis of the Jurassic succession of the Sverdrup Basin, Canadian Arctic Archipelago. *Canadian Journal of Earth Sciences*, 30, 301-320.

Embry, A.F. 1997. Global sequence boundaries of the Triassic and their recognition in the Western Canada Sedimentary Basin. *Bulletin Canadian Petroleum Geology*, 45, 415 – 433.

Embry, A., 2011, Petroleum Prospectivity of the Triassic-Jurassic Succession of Sverdrup Basin, Canadian Arctic Archipelago. *In*: Spencer, A. M., Embry, A. F., Gautier, D. L., Stoupakova, A. V. & Sørensen, K. (eds) Arctic Petroleum Geology. Geological Society, London, Memoirs, 35, 545–558.

Embry, A. and Beauchamp, B. 2008. Sverdrup Basin. *In*: Miall, A. (ed.) Sedimentary Basins of the World, Vol. 5; The Sedimentary Basins of the United States and Canada. Elsevier, Amsterdam, 451–472.

- Fritz, W. et al. 1992, Cambrian to Middle Devonian Assemblages. In: Gabrielse, H. and Yorath, C. (eds), *Geology of the Cordilleran Orogen*, Geological Survey of Canada, *Geology of Canada* No. 4, p.151-217.
- Gradstein, F., Ogg, J., Schmitz, M. and Ogg, G., 2012, *The Geologic Time Scale 2012*. Elsevier, Amsterdam, 1043 p.
- Harrison, J.C. 1995. Melville Island's salt-based fold belt, Arctic Canada. Geological Survey of Canada, *Bulletin* 472, 331 p.
- Harrison, C. et al. 1999, Correlation of Cenozoic sequences of the Canadian Arctic region and Greenland; implications for the tectonic history of northern North America. *BCPG*, V.47, p.223-254.
- Helwig, J., Kumar, N., Emmet, P. and Dinkelman, M. G. Regional seismic interpretation of crustal framework, Canadian Arctic passive margin, Beaufort Sea, with comments on petroleum potential. In: Spencer, A. M., Embry, A. F., Gautier, D. L., Stoupakova, A. V. and Sørensen, K. (eds) *Arctic Petroleum Geology*. Geological Society, London, *Memoirs*, 35, 527-544.
- Higgins, A.K., Ineson, J.R., Peel, J.S., Surlyk, F. and Scnderholm, M. 1991a. Cambrian to Silurian basin development and sedimentation, North Greenland. In: H.P. Trettin (ed.). *Geology of the Innuitian Orogen and Arctic Platform of Canada and Greenland*. Geological Survey of Canada, *Geology of Canada*, no. 3, p. 111-161.
- Houseknecht, D.W., and Bird, K.J., 2004, Sequence stratigraphy of the Kingak Shale (Jurassic – Lower Cretaceous), National Petroleum Reserve in Alaska: *American Association of Petroleum Geologists Bulletin*, v. 88, p. 279–302.
- Mayr, U., 1978, Stratigraphy and correlation of lower Paleozoic formations, subsurface of Cornwallis, Devon, Somerset, and Russell islands, Canadian Arctic Archipelago: Geological Survey of Canada *Bulletin* 276, 55 p.
- Mayr, U., J. J. Packard, Q. H. Goodbody, A. V. Okulitch, R. J. Rice, F. Goodarzi, and K. R. Stewart, 1994, The Phanerozoic geology of southern Ellesmere and northern Kent islands, Canadian Arctic Archipelago: Geological Survey of Canada *Bulletin* 470, 298 p.
- Mayr, U., T. De Freitas, B. Beauchamp, and G. Eisbacher, 1998, The geology of Devon Island north of 76, Canadian Arctic Archipelago: Geological Survey of Canada *Bulletin* 526, 500 p.
- Poulton, T. P., Leskiw, K., and Audretsch, A. P., 1982, Stratigraphy and microfossils of Jurassic Bug Creek Group of northern Richardson Mountains, northern Yukon and adjacent Northwest Territories, Geological Survey of Canada, Ottawa, Canada, *Bulletin* 325, 130 p.
- Pyle, L. and Jones, A. 2009, Regional Geoscience Studies and Petroleum Potential of Peel Plateau and Plain, NWT and YT, Project Volume. NWT Open File 2009-02, 549 p.
- Ricketts, B. 1991. Delta evolution in the Eureka Sound Group, western Axel Heiberg Island: the transition from wave-dominated to fluvial-dominated deltas; Geological Survey of Canada, *Bulletin* 402, 72 p.
- Ricketts, B. 1994. Basin analysis, Eureka Sound Group, Axel Heiberg and Ellesmere islands, Canadian Arctic Archipelago. Geological Survey of Canada, *Memoir* 439, 119 p.
- Ricketts, B. and Stephenson, R., 1994, The Demise of Sverdrup Basin: Late Cretaceous-Paleogene Sequence Stratigraphy and Forward Modeling. *Journal of Sedimentary Research*, v.64, p. 516-530.
- Rolf, T, Coltice, N. and Tackley, P., 2012, Linking continental drift, plate tectonics and the thermal state of the Earth's mantle. *Earth and Planetary Science Letters*, v. 351, p. 134-146.
- Saltzman, M., and Thomas, E. 2012, Carbon Isotope Stratigraphy. In: Gradstein, F., Ogg, J., Schmitz, M. and Ogg, G., (eds.), *The Geologic Time Scale 2012*. Elsevier, Amsterdam, p. 207-232.
- Schenk, C. J. Geology and petroleum potential of the West Greenland–East Canada Province. In: Spencer, A. M., Embry, A. F., Gautier, D. L., Stoupakova, A. V. and Sørensen, K. (eds) *Arctic Petroleum Geology*. Geological Society, London, *Memoirs*, 35, 627-646.
- Sloss, L. L., 1988, Tectonic evolution of the craton in Phanerozoic time, in Sloss, L. L. ed., *Sedimentary cover—North American craton: U.S.*, The Geology of North America, Geological Society of America, Boulder, CO, v. D-2, pp. 25–51.
- Thorsteinsson, R., and U. Mayr, 1987, The sedimentary rocks of Devon Island, Canadian Arctic Archipelago: Geological Survey of Canada *Memoir* 411, 182 p.
- Trettin, H.P. Mayr, U., Long, G.D.F. and Packard, J.J. 1991. Cambrian to Early Devonian Basin Development, Sedimentation and Volcanism, Arctic Islands. In: H.P. Trettin (ed.). *Geology of the Innuitian Orogen and Arctic Platform of Canada and Greenland*. Geological Survey of Canada, *Geology of Canada*, no. 3, p. 165-238.