

Dolomitization of the Presqu'ile Barrier: An Alternative Paleohydrological Model

Mario Coniglio¹, Dave Morrow², and Nick Wilson³

¹University of Waterloo, Waterloo, Ontario, Canada (coniglio@uwaterloo.ca)

²Geological Survey of Canada, Calgary, Alberta, Canada

³Talisman Energy Inc., Calgary, Alberta, Canada

Abstract

Regional geographic trends of dolomite stable oxygen isotope values, radiogenic strontium ratios and fluid inclusion homogenization temperatures were interpreted in earlier studies of the Presqu'ile barrier to reflect a basin-scale, eastward up-dip migration of hot, radiogenic fluids driven by tectonic compression, sedimentary loading and uplift of the western part of the Western Canada Sedimentary Basin. These dolomitizing fluids were thought to have been derived by a combination of topographic recharge and tectonic compaction during either the Cretaceous- Tertiary Laramide Orogeny (probably the main phase) or possibly the earlier Antler Orogeny (Devonian-Carboniferous). Recent reexamination of the petrography and geochemistry of the hydrothermal, fabric-destructive dolomites from 4 diamond drill cores from the Pine Point Mines property and 31 petroleum cores from the subsurface of the NWT to the west of Pine Point (Fig. 1), and reconsideration of the previously derived data, provide a basis for challenging this model. Overall, newly obtained geochemical and fluid inclusion data are remarkably variable, and show inconsistent relationships with both petrographic and cathodoluminescence characteristics.





Figure 1. Locations of wells sampled and from previous studies (Qing and Mountjoy 1994) and locations of major faults in the study area.

Figure 2 illustrates the non-correspondence between oxygen isotopes from samples of Presqu'île Dolomite and their geographic location



deep topographic recharge or by “Tectonic” compaction-driven fluid flow. Variability in the isotopic data, particularly stable oxygen isotope values, even at a borehole scale, can be explained by temporal or spatial variations in heat flow and/or depth to basement, or to the degree of recrystallization as a function of fluid-rock interaction.

References Cited

Janicki, E.P., 2004, Selected Lithologs of Presqu'ile Dolomite in the southeastern Great Slave Plain, Northwest Territories: C.S. Lord Northern Geoscience Centre, Yellowknife, Northwest Territories, NWT Open Report 2004-001.

Qing, H., and E.W. Mountjoy, 1994, Formation of coarsely-crystalline, hydrothermal dolomite reservoirs in the Presqu'ile barrier, Western Canada Sedimentary Basin: AAPG Bulletin, v. 78, p. 55-77.