

## Overpressure, Seal Integrity, and Reservoir Connectivity in the Scotian Basin

Carla H. Dickson<sup>1</sup>, Ricardo L. Silva<sup>1</sup>, Grant Wach<sup>1</sup>, and Neil Watson<sup>2</sup>

<sup>1</sup>Basin and Reservoir Lab, Department of Earth Sciences, Dalhousie University, Halifax, Nova Scotia, Canada ([carla.h.dickson@dal.ca](mailto:carla.h.dickson@dal.ca))

<sup>2</sup>Atlantic Petrophysics Limited, Halifax, Nova Scotia, Canada ([neil.watson@huskyenergy.com](mailto:neil.watson@huskyenergy.com))

### Abstract

Overpressure is abnormally high subsurface pressure that exceeds hydrostatic pressure at a given depth, and occurs when connate fluids are unable to escape the surrounding mineral matrix. Causes for overpressure include the compaction effect (under compaction), formation foreshortening, faulting, massive evaporite deposition, mineral phase change, repressuring, and hydrocarbon generation. Often obstructed fluid migration pathways are also a factor, and more than one of the above reasons contributes to the formation of overpressure in a system.

To maintain overpressure within a reservoir, compartmentalization is required, where a petroleum accumulation is separated into several discrete fluid/pressure compartments, which occurs when hydrocarbon migration is prevented across sealed boundaries within the reservoir. The boundaries can be divided into two groups: static seals that completely seal and are able to trap hydrocarbons for a geologic time-frame, and dynamic seals that are low permeability baffles to flow that decrease rate of flow to essentially zero at the production scale but will allow for eventual equalization over a geologic scale.

Overpressured sediments tend to exhibit better porosity than would be normally be predicted from their depth, therefore make attractive prospects for hydrocarbon reservoirs. Difficulty managing abnormal pressures during drilling is frequently cited for abandoning wells or for unexpected, expensive changes in the casing program. The causes of overpressure in the Scotian Basin are poorly understood, and with the recent increase in exploration for hydrocarbons offshore Nova Scotia, understanding the development of overpressure is important for maximizing safety, production, and developing new opportunities.

The Scotian Basin comprises Mesozoic and Cenozoic sediments overlying a Paleozoic basement, creating a structurally complex basin containing carbonate banks, platforms, and ridges within a set of differentiated sub-basins. There has been over 250 Ma of continuous sedimentation to the basin, predominantly sourced from the Appalachian Orogen then transported by the paleo-St. Lawrence River. The Mesozoic-Cenozoic infill of the basin reaches a maximum thickness of 16km (Wade, MacLean, and Williams, 1995). Several intervals have potential as reservoirs (e.g. Eurydice, Mississauga and MicMac formations) with several of these in production.

The petroleum system of the Sable Subbasin as identified by Grant et al. (1984) showed that hydrocarbon accumulations were prone to gas, small volumes, and scattered both vertically and laterally. Richards et al. (2008) identified that reservoirs in the Scotian Basin are often connected, and where reservoirs are thick and well developed there tends to be an absence of sufficient seals. This study investigated hydrocarbon distribution, resource potential, and production performance of the Sable Subbasin and included structural geometrical controls on

the fluid distribution and pressures in the region. They suggested overpressure on the Scotian Basin might be associated with particular trap styles.

The reservoir connectivity analysis is important to understanding the overpressure systems of the Scotian Basin, and includes reservoir architecture, definition of seal integrity, and determination of connectedness for compartments. The use of analogous basins with similar formation and depositional histories is also essential, as data available from the Scotian Basin is limited; other systems on the Atlantic conjugate margin include Jeanne d'Arc Basin, Porcupine Basin, Northern North Sea Basin, and the Eastern Venezuela Basin. Overpressure is a risk element in all of these basins including the Scotian Basin, and needs to be understood to safely develop and produce offshore resources.

### References Cited

Grant, A., McAlpine, K., and Wade, J. (1984). The Geological Margin of Eastern Canada: Geological Framework and Petroleum Potential, in Halbouty MT (ed.). *Future Petroleum Provinces of the World: AAPG Bulletin*, 177-205.

Madatov, A. (2005a). The overpressure driven seismic velocity response. Review of standard models and methods for extraction in the context of basin modelling approach to overpressure prediction. *Proceedings of the Murmansk State Technical University*, 8(1), 84-119.

National Research Council of Canada. (2010). NRC National Science Library. Retrieved April 2013, from <http://cisti-icist.nrc-cnrc.gc.ca/eng/ibp/cisti/index.html>.

Richards, B., Fairchild, L., Vrolijk, P., & Hippler, S. (2008). Reservoir Connectivity Analysis, Hydrocarbon Distribution, Resource Potential, and production Performance in the Clastic Plays of the Sable Subbasin, Scotian Shelf. *Central Atlantic Conjugate Margins Conference*, 165-185.

Wade, J., MacLean, B., and Williams, G. (1995). Mesozoic and Cenozoic stratigraphy, eastern Scotian Shelf: new interpretations. *Canadian Journal of Earth Sciences*, 32(9), 1462-1473.

Webster, M., O'Connor, S., Pindar, B., and Swarbrick, R. (2011, March). Overpressures in the Taranaki Basin: Distribution, causes, and implications for exploration. *AAPG Bulletin*, 95(3), 339-370.

Wielens, J. (2003). Overpressures on the Scotian Shelf. Open File Report 1557. Geological Survey of Canada.

Yassir, N., and Bell, J. (1994). Relationship between pore pressure, stresses, and present-day geodynamics in the Scotian Shelf, offshore Eastern Canada. *AAPG Bulletin*, 78(12), 1863-1880.