Tidally-Influenced Sedimentation on a High-Latitude Cretaceous Coastal Plain: The Prince Creek Formation, North Slope, Alaska*

Peter P. Flaig¹, Paul J. McCarthy², and Anthony R. Fiorillo³

Search and Discovery Article #50365 (2010)
Posted December 17, 2010

*Adapted from oral presentation at AAPG International Conference and Exhibition, Calgary, Alberta, Canada, September 12-15, 2010

Abstract

The late Cretaceous (Early Maastrichtian) Prince Creek Formation on Alaska's North Slope records deposition on a high-latitude coastal plain within tidally-influenced first-order meandering trunk channels, second-order meandering distributary channels, third-order fixed (anastomosed?) distributary channels and vegetated floodplains. Regionally restricted suspended-load meandering trunk channels (13-17 m thick) include a basal conglomerate, a medium-grained multi-storey sandbody lacking roots that fines-upward into inclined heterolithic stratification (IHS) and 3-4 m of finer organic-rich sediments. Meandering distributaries (2-6 m thick) composed of a singlestorey, fine-grained, rooted sandbody dominated by IHS and including abundant mud-filled channel plugs occur as laterally-extensive heterolithic sheets encased in organic-rich floodplain facies that appear lenticular along strike. Fixed (anastomosed?) distributary channels are composed of a fine-grained, single-storey, current rippled and rooted sandbody atop an arcuate erosive base with rare IHS and siltstone "wings" that extend laterally from the uppermost channelfill. Numerous fixed (anastomosed?) channels occupy tiers at the same stratigraphic level, with fixed channels incised into meandering distributaries or floodplain facies.

All channels are interbedded with drab-colored, organic-rich floodplain facies interpreted as the deposits of levees, splays, lakes, ponds, swamps, marshes, mires, paleosols, and ashfall. Spatial relationships between channel types, and between channels and floodplains indicate that most deposition took place adjacent to trunk channels on crevasse splay-complexes separated by organic-rich

¹Jackson School of Geosciences, Bureau of Economic Geology, University of Texas at Austin, Austin, TX (peter.flaig@beg.utexas.edu)

²Dept. of Geology and Geophysics, and Geophysical Institute, University of Alaska Fairbanks, Fairbanks, AK

³Museum of Nature and Science, Dallas, TX

floodplain facies. The lateral migration of meandering distributaries driven by peat compaction and higher flow velocities and the vertical filling of fixed (anastomosed?) distributaries may have been contemporaneous. Alternatively fixed (anastomosed?) distributaries may record the initial or waning stages of flow during splay-complex formation or abandonment. IHS composed of rhythmically-repeating, coarse-to-fine couplets of current rippled sandstone/siltstone and mudstone is found within all channel-forms. The rhythmic and repetitive nature of IHS couplets together with relatively thick, muddy fine-grained members in the IHS suggest that flow within channels was likely influenced by tidal effects. Locally abundant jarosite lends credence to an interpretation of marine-influenced sedimentation.

References

Bird, K.J., 1987, North Slope oil and gas; the Barrow Arch paradox: SEPM Field Trip Guidebook, Pacific Section, v. 50, p. 41.


Bird, K.J. and C.M. Molenaar, 1992, The North Slope foreland basin, Alaska: AAPG Memoir, v. 55, p. 363-393.

Decker, D., J.E. Box, and R. Benson, 2009, Greenland ice sheet tidewater glacier front changes; seasonal and interannual changes 2000-2008: Association of American Geographers, Annual Meeting Abstracts, p. 149.


Fiorillo, A.R. and R. Gangloff, 2000, Theropod teeth from Prince Creek Formation (Cretaceous) of Northern Alaska, with speculations on Arctic dinosaur paleoecology: Arctic Science Conference, Program and Abstracts, v. 51, p. 87.

Garrity, C.P., D.W. Houseknecht, K.J. Bird, C.J. Potter, T.E. Moore, P.H. Nelson, and C.J. Schenk, 2005, USUS 2005 oil and gas resource assessment of the central North Slope, Alaska; play maps and results: USGS Open File Report #2005-1182,


Mull, C.G., D.W. Houseknecht, and K.J. Bird, 2003, Revised Cretaceous and Tertiary stratigraphic nomenclature in the Colville Basin, northern Alaska: USGS Professional Paper 1673, version 1.0. http://pubs.usgs.gov/pp/p1673/index.html




Purpose of this talk



Location:



Regional Stratigraphy:



Prince Creek Formation records deposition in a foreland basin on a high-latitude coastal plain

Colville Basin...Why Important?

- Preserves a high-latitude dinosaur ecosystem
- Surface expression of subsurface reservoirs





FIGURE 3. Schematic structural and stratigraphic cross section across the central North Slope (from Bird, 1987). Shaded zone between Torok Formation and Kingak Shale is the pebble shale unit and gamma-ray zone. See Figure 2 for location of section.

BIRD & MOLENAAR, 1992

Study Area 75 measured stratigraphic sections

Grain-size, lithology, contacts, sed. structures, flora, fauna



151° 30' W

Sand clmdsivffmcg

FA-IV


FA-IV


C m

C m


A De

Typical Facies







Typical Facies



Facies Associations



FA-IV Interbedded Sandstone and Siltstone



151° 30' W **Paleoflow** Ocean **Point** 70° 00' N Vector mean=60° **68** Paleoflow orientation Number of measurements




Axial flow recorded from trough cross-laminations

(consistent with previous work)



Large Sinuous Channels (FA-I)





Only Channels with Gravels/Boulders




...some clasts are the size a football ...important why?...this is <u>extremely</u> rare

Inclined Heterolithic Stratification (IHS)





Small Sinuous Channels (FA-II)



- Fine grained sand and silt
- Alternating sand & mud (couplets)
- Carbonaceous roots throughout
- Small sinuous channels=

 <u>Meandering second-order</u>

 <u>distributary channels</u>






4.5 m

Opposing apparent dips


Small Low Sinuosity Channels (FA-III)




Small low-sinuosity channels= Third order fixed (anastomosed?) distributaries



All Channels have rhythmically-repeating sand/mud couplets = Tidal Influence




Facies Associations




Levees





Small Floodplain Lakes





Small-scale Crevasse Splays Jarosite Halos sandstone CUS siltstone mudstone shale-Jarosite


Dinosaur Bone


















Raleosol Microfacies















Comparison with Modern Soils




Jarosite mottling in acid sulphate soil (Mitch Tulau)

Estuarine soil in Australia


Modern Inceptisols/Entisols/Potential Acid
Sulphate Soils

Alluvial Architecture

Large Sinuous Channels (FA-I)



Rare and regionally-restricted




Most common channel-form

Small Sinuous Channels (FA-II)



Deposits appear lenticular along strike

Small Low Sinuosity Channels (FA-III)



Incise into meandering distributaries or floodplains

Interbedded Sandstone, Siltstone, Mudstone (FA-IV)

Floodplain

Thin Sheet Sands 30 m Carbonaceous shale and coal


Depositional Model

Trunk Channels

Distributary Channels on Crevasse Splay

Complexes

Organic-rich floodplains



Conclusions:

- Channels
 - Larger first-order meandering trunk channels (rare)
 - Smaller second-order meandering distributary channels
 - Small third-order anastomosed(?) distributary channels
- IHS (rhythmically-repeating sand/mud couplets) in channels suggest tidal influence on system (abundant jarosite, pyrite, and brackish-water fauna also indicate proximity to marine waters)
- Floodplain Depositional Environments:
 - Small-scale crevasse splays
 - Levees
 - Lakes, ponds, swamps, ashfall deposits
 - Paleosols

Conclusions: Paleosols and Pollen

-Paleosols:

- Weakly-developed, cumulative to compound soils
- Frequent influx of material into soils
- Predominantly waterlogged
- Drying-out of parts of the floodplain
- Marine influence
- Similar to modern Entisols, Inceptisols, and Potential Acid Sulphate soils
- Depositional Model
 - -Meandering trunk channels with distributary channels forming <u>crevasse-splay complexes</u> that alternate with organic floodplain facies

Thank You!







Three Field Seasons 30+ individuals Logistical Support









NSF grants: OPP-0425636 to Paul McCarthy OPP-424594 to Anthony Fiorillo Also David LePain, Paul Layer

The Geist Fund at the University of Alaska Museum of the North

The Alaska Geological Society

The Geological Society of America

The Evolving Earth Foundation

The University of Alaska Graduate School Fellowship Program

British Petroleum

Dolores van der Kolk



