

Fifty Shades of Grey: Utilizing “Conventional” Sedimentology and Sequence Stratigraphy to Unlock Rock Quality to Reservoir Quality Relationships in the Liquids Rich Duvernay Shale Play, Kaybob Alberta Canada*

Lindsay Dunn¹, Keith Gordon¹, and Marc Houle¹

Search and Discovery Article #10702 (2015)

Posted January 19, 2015

*Adapted from extended abstract prepared in conjunction with presentation at CSPG/CSEG/CWLS GeoConvention 2013, (Integration: Geoscience engineering Partnership) Calgary TELUS Convention Centre & ERCB Core Research Centre, Calgary, AB, Canada, 6-12 May 2013, Datapages/CSPG © 2015

¹Athabasca Oil Corp., Calgary, Alberta, Canada (ldunn@atha.com)

Abstract

The Duvernay Formation of Alberta is a mudrock reservoir that produces oil and gas from mature to overmature oil-prone source rocks. What makes the Duvernay Formation particularly attractive as a “shale” play like its American cousin the Eagle Ford Formation is its ability to produce significant volumes of liquids due to its over pressured nature. The highest reservoir quality within the Duvernay is developed in high total organic carbon (TOC), siliceous mudstones. A strong positive correlation between silica content and TOC in the siliceous mudstones indicates a biogenic rather than detrital source for much of the silica and produces a higher modulus, brittle rock amenable to fracture stimulation (Dunn et al., 2012). A strong positive correlation between increased reservoir quality and TOC indicates an organo-porosity system analogous to the Barnett and Marcellus (Dunn et al., 2012).

Introduction

Contrary to many people’s perception, these “basinal” source rock deposits are inherently heterogeneous due to the dynamic nature of their depositional environment. Variations in primary productivity, bottom water energy, rates/process of sediment accumulation and redox conditions, result in a complex depositional system (May and Anderson, 2011). Relative amounts of TOC, biogenic silica, clay, and carbonate control reservoir quality in the Duvernay vary systematically. “Conventional” sedimentology and sequence stratigraphic workflows provide the framework to develop a predictive model deciphering rock quality to reservoir quality relationships both vertically and laterally.

Discussion

In the study area, the Duvernay is underlain by the Majeau Lake Formation and overlain by the Ireton Formation representing a single second order depositional cycle (Switzer et al., 1994). The Duvernay Formation deposited during the late Transgressive Systems Tract (TST) early Regressive Systems Tract (RST). High-resolution macroscopic core descriptions are integrated with lab analyses and petrophysics to develop a higher order sequence stratigraphic framework in the Kaybob Study area.

The underlying Majeau Lake Formation and Duvernay Formation succession comprises at least four 3rd order transgressive-regressive cycles (T-R) superimposed on the second order systems tract ([Figure 1](#)). Within an individual T-R cycle, the mudstone becomes increasingly rich in organics and biogenic silica through the transgressive systems tract and progressively diluted by extra-basinal clay and carbonate detritus through the overlying regressive systems tract.

Summary

This sequence stratigraphic model has been integrated with the Duvernay Depositional Model ([Figure 2](#)) and can be used not only to high grade geographic areas but can also be used to pick optimum horizontal drilling targets and/or completion zones in order to maximize production rates.

Acknowledgements

Thank you to Athabasca Oil Corp. for the permission to present this ongoing work.

References Cited

Dunn, L.A., G. Schmidt, K. Hammermaster, M. Brown, R. Bernard, E. Wen, R. Befus, and S. Gardiner, 2012, The Duvernay Formation (Devonian): Sedimentology and Reservoir Characterization of a Shale Gas/Liquids play in Alberta, Canada: Canadian Society of Petroleum Geologists Annual Core Convention program with abstracts, May 2012.

May, J.A., and D.S. Anderson, 2011, Mudrock Reservoirs – Why Depositional Fabric and Sequence Stratigraphic Framework Matter: AAPG Search and Discovery Article #90124, Web accessed January 3, 2015, http://www.searchanddiscovery.com/pdfz/documents/2013/80338may/ndx_may.pdf.html.

Stoakes, F.A., 1980, Nature and Control of Shale Basin Fill and its Effect on Reef Growth and Termination: Upper Devonian Duvernay and Ireton Formations of Alberta, Canada: Bulletin of Canadian Petroleum Geology, v. 28/3, p. 345-410.

Switzer, S.B., W.G. Holland, D.S. Christie, G.C. Graf, A. Hedinger, R. McAuley, R. Wierzbicki, and J.J. Packard, 1994, Devonian Woodbend - Winterburn strata of the Western Canada Sedimentary Basin: in G.D. Mossop and I. Shetsen (eds.), Geological Atlas of the Western Canada Sedimentary Basin, Calgary, Canadian Society of Petroleum Geologists and the Alberta Research Council, Chapter 12.




Figure 1. Proposed sequence stratigraphic model for the Duvernay Formation Kaybob, Alberta Canada.




Figure 2. Integrated Duvernay depositional and sequence stratigraphic model for the Kaybob Area. Axial transport of clastic detritus material and estimated water depths as per Stoakes (1980).